



Rapid QSM Acquisition with Wave-CAIPI

**B. Bilgic^{1,2}, B. A. Gagoski^{2,3}, S.F. Cauley^{1,2}, A.P. Fan^{1,4},
J.R. Polimeni^{1,2}, P.E. Grant^{2,3}, L.L.Wald^{1,2,5}, K. Setsompop^{1,2}**

1 Martinos Center for Biomedical Imaging, Charlestown, MA,

2 Harvard Medical School, Boston, MA


3 Boston Children's Hospital, Boston, MA

4 Massachusetts Institute of Technology, Cambridge, MA

5 Harvard-MIT Division of Health Sciences and Technology, Cambridge, MA



Highly Accelerated 3D Cartesian Imaging

- Recent modifications to rectilinear k-space sampling provided more robust reconstructions of highly under-sampled datasets





Highly Accelerated 3D Cartesian Imaging

- Recent modifications to rectilinear k-space sampling provided more robust reconstructions of highly under-sampled datasets




Highly Accelerated 3D Cartesian Imaging

- Recent modifications to rectilinear k-space sampling provided more robust reconstructions of highly under-sampled datasets




Highly Accelerated 3D Cartesian Imaging

- Recent modifications to rectilinear k-space sampling provided more robust reconstructions of highly under-sampled datasets




Effect of G_y in image space




Wave-CAIPI Sampling

- Recent modifications to rectilinear k-space sampling provided more robust reconstructions of highly under-sampled datasets




- Wave-CAIPI: 2D CAIPI + BPE in 2 directions
- Spread aliasing in 3D to take full advantage of 3D coil profiles




Wave-CAIPI causes voxel spreading in 3 dimensions

- Combination of G_y and G_z gradients with inter-slice shifts yields voxel spreading across three dimensions




Wave-CAIPI causes voxel spreading in 3 dimensions

- Combination of G_y and G_z gradients with inter-slice shifts yields voxel spreading across three dimensions




Wave-CAIPI causes voxel spreading in 3 dimensions

- Combination of G_y and G_z gradients with inter-slice shifts yields voxel spreading across three dimensions




Wave-CAIPI causes voxel spreading in 3 dimensions

- Combination of G_y and G_z gradients with inter-slice shifts yields voxel spreading across three dimensions




Wave-CAIPI improves parallel imaging

- Voxel spreading increases the distance across aliasing locations
- This increases the variation in coil sensitivity profiles and improves parallel imaging capability




R=3x3 @ 3 Tesla, 1 mm iso, $T_{acq}=2.3\text{min}$

k-space




Recon




2D-CAIPI

1/G-factor




R=3x3 @ 3 Tesla, 1 mm iso, $T_{acq}=2.3\text{min}$

k-space





Recon



2D-CAIPI



1/G-factor




Bunch Encoding




R=3x3 @ 3 Tesla, 1 mm iso, $T_{acq}=2.3\text{min}$

k-space





Recon






2D-CAPI



1/G-factor



Bunch Encoding




Wave-CAPI




R=3x3 @ 7 Tesla, 1 mm iso, $T_{acq}=2.3\text{min}$

k-space




Recon




2D-CAIPI

1/G-factor




R=3x3 @ 7 Tesla, 1 mm iso, $T_{acq}=2.3\text{min}$

k-space




Recon

2D-CAIPI





1/G-factor


$G_{max}=1.74$



Bunch Encoding




$G_{max}=2.12$




R=3x3 @ 7 Tesla, 1 mm iso, $T_{acq}=2.3\text{min}$

k-space





Recon




2D-CAPI




1/G-factor



Bunch Encoding



Wave-CAPI



R=3x3 @ 7 Tesla, 1 mm iso, $T_{acq}=2.3\text{min}$

Wave-CAIPI

